

# **Intermediate Lake Shoreline Survey 2016**

*By Tip of the Mitt Watershed Council*

Report written by:

**Matt L. Claucherty**

*Monitoring and Research Coordinator*

## Table of Contents

	Page
<b>List of Tables and Figures</b> .....	<b>iii</b>
<b>Summary</b> .....	<b>2</b>
<b>Introduction</b> .....	<b>3</b>
Background.....	3
Shoreline Development Impacts.....	4
Study Area.....	7
<b>Methods</b> .....	<b>14</b>
Field Survey Parameters.....	14
Data Processing.....	17
<b>Results</b> .....	<b>18</b>
<b>Discussion</b> .....	<b>24</b>
Recommendations.....	28
<b>Literature and Data Referenced</b> .....	<b>30</b>

## List of Tables

	<b>Page</b>
Table 1. Intermediate Lake Watershed land-cover statistics.....	8
Table 2. Categorization system for <i>Cladophora</i> density.....	15
Table 3. <i>Cladophora</i> density results.....	18
Table 4. Greenbelt rating results.....	20
Table 5. Shoreline alteration results.....	22
Table 6. Shoreline erosion results.....	22
Table 7. Critical shoreline survey parameter comparisons: 2000 to 2016.....	25
Table 8. Shore survey statistics from Northern Michigan lakes.....	26

## List of Figures

	<b>Page</b>
Figure 1. Map of Intermediate Lake and Watershed.....	10
Figure 2. Chart of average Secchi depths in Intermediate Lake.....	11
Figure 3. Chart of chlorophyll-a concentrations in Intermediate Lake.....	12
Figure 4. Total phosphorus trends in Intermediate Lake 1995 to 2016 (TOMWC 2016).....	13
Figure 5. <i>Cladophora</i> algae density results for Intermediate Lake.....	19
Figure 6. Greenbelt score totals results for Intermediate Lake.....	21
Figure 7. Shoreline erosion severity results for Intermediate Lake.....	23

## SUMMARY

During the early summer of 2016, the Tip of the Mitt Watershed Council conducted a shoreline survey of Intermediate Lake as part of a comprehensive shoreline survey for the entire Elk River Chain of Lakes. Watershed Council staff and interns surveyed the upper Elk River Chain (Beals Lake through Intermediate Lake) in 2016 and the lower Elk River Chain (Lake Bellaire through Elk Lake) in 2017. Surveys were designed to document conditions that can potentially impact water quality, including the three biggest threats to inland lakes: nutrient pollution, habitat loss, and shoreline erosion.

Survey results indicate that human activity along the Intermediate Lake shoreline is likely impacting the lake ecosystem and water quality. Nearly two thirds (63%) of all shoreline properties had little to no vegetation growing at water's edge. Vegetation removal is concentrated in the northern portion of the lake, especially along the western shoreline. The southern basin, flanked by state land, provides the greatest amount of natural shoreline habitat. Erosion, found at 53% of all properties, poses a threat to the Lake's ecosystem and aesthetic quality.

On a positive note, signs of nutrient pollution are not widespread. *Cladophora*, an algal indicator of nutrient pollution, was documented at 19% of all properties, most of which was light or very light in density. This suggests that failing septic systems and heavy lawn fertilization issues are not occurring as commonly in Intermediate Lake as in other Northern Michigan Lakes.

Steps can be taken to improve the habitat and water quality of Intermediate Lake. Erosion sites can be repaired, vegetation can be allowed to regrow on the shoreline, providing improved pollutant filtration and erosion resistance. Outreach to shoreline property owners regarding lake-friendly shoreline management practices can help to improve conditions. Educating residents on beneficial and harmful activities is often all that is needed to bring about change.

## INTRODUCTION

### **Background:**

During the late spring of 2016, a shoreline survey was conducted on Intermediate Lake by the Tip of the Mitt Watershed Council (Watershed Council) to document shoreline conditions that potentially impact water quality. The entire shoreline was surveyed to document the following: algae as a nutrient pollution indicator, erosion, shoreline alterations, greenbelts, and tributary inlets and outlets. This survey was funded by a grant from the Michigan Department of Environmental Quality as part of a larger Elk River Watershed protection initiative.

In 2000, Clean Water Act funds were granted to eight organizations and agencies by the Michigan Department of Natural Resources for a three year project to protect water resources of the Elk River/Chain of Lakes Watershed. One of the project's tasks was a shoreline water resource/nonpoint source pollution survey. The survey was intended to document basic information for educational purposes and as a basis for water resource management decisions.

The 2000 survey was primarily conducted by small boat traveling closely along the shoreline. The description of shoreline property parcels, as observed from the water, was matched with maps showing parcel lines, tax numbers, and ownership information obtained from the equalization departments of the four counties covered by the watershed. This allowed the location of observed resource features to be later plotted on the maps with relative accuracy. The survey results were entered into a computerized database and depicted on a map generated by a geographic information system. In 2000, shoreline survey efforts documented shoreline erosion and *Cladophora* algae, but did not document vegetation removal or shoreline alterations.

The 2016 survey provides a comprehensive data set documenting shoreline conditions on Intermediate Lake; a valuable data set that can be used as a lake management tool. Combined with follow-up activities, such as questionnaires and on-site visits, problems in shoreline areas

that threaten the lake's water quality can be identified and corrected. These solutions are often simple and low cost, such as regular septic system maintenance, proper lawn care practices, and wise land use along the shoreline. Prevention of problem situations can also be achieved through the publicity and education associated with the survey. Periodic repetition of shoreline surveys is important for identifying new and chronic problem sites, determining long-term trends of near-shore nutrient inputs and shoreline alterations associated with land-use changes, and for assessing the success of remedial actions.

### **Shoreline Development Impacts:**

Lake shorelines are the critical interface between land and water; where human activity has the greatest potential for degrading water quality. Traditional development of shoreline properties for residential, commercial, or other uses invariably leads to negative impacts on the lake ecosystem. During the development process, the natural landscape is altered in a variety of ways: vegetation is removed, the terrain is graded, utilities are installed, structures are built, and areas are paved. These changes to the landscape and subsequent human activity in the shoreline area have consequences on the aquatic ecosystem. Nutrients from organic wastes, contaminants from cars and roads, and soils from eroded areas are among some of the pollutants that end up in and negatively impact the lake following shoreline development.

Nutrient pollution can have adverse impacts on aquatic ecosystems and pose a danger to human health. While nutrients are necessary to sustain a healthy aquatic ecosystem, excess nutrients will stimulate nuisance aquatic plant growth of both macrophytes (aquatic plants that grow in or near water and are either emergent, submergent, or floating) and algae.

Additionally, algal blooms pose a public health risk as some species (i.e. blue green algae) produce toxins, including hepatotoxins (toxins that cause liver damage) and neurotoxins (toxins that affect the nervous system). Excess plant and algae growth can also degrade water quality by depleting the ecosystem's dissolved oxygen stores. During nighttime respiration, plants compete with other organisms for a limited oxygen supply. Furthermore, the decomposition of algae and plants has the potential to deplete dissolved oxygen supplies due to the aerobic

activity of decomposers, particularly in the deeper waters of stratified lakes.

Surface waters receive nutrients through a variety of natural and cultural (human) sources. Natural sources of nutrients include stream inflows, groundwater inputs, surface runoff, organic inputs from riparian (shoreline) areas, and atmospheric deposition. Springs, seeps, streams, and artesian wells are often naturally high in nutrients due to the geologic strata they encounter. Nearby wetland seepages may also discharge nutrients at certain times of the year. Cultural sources include septic systems, fertilizers, and stormwater runoff from roads, driveways, parking lots, roofs, and other impervious surfaces. Poor agricultural and forestry practices, which oftentimes result in soil erosion, and wetland destruction also contribute to nutrient pollution. Furthermore, some cultural sources (e.g., malfunctioning septic systems) pose a potential health risk due to bacterial and viral contamination.

Severe nutrient pollution is detectable through chemical analyses of water samples, physical water measurements, and the utilization of biological indicators (a.k.a., bio-indicators). Although chemical analyses of water samples to check for nutrient pollution can be effective, they are oftentimes more labor intensive and costlier than other methods. Typically, water samples are analyzed to determine nutrient concentrations (usually forms of phosphorus and nitrogen), but other chemical constituents, such as chloride, can be measured. Physical measurements, such as water temperature and conductivity (i.e., the water's ability to conduct an electric current), are primarily used to detect malfunctioning septic systems. Biologically, nutrient pollution can be detected along the lake shore by noting the presence of *Cladophora* algae, a bio-indicator.

*Cladophora* is a branched, filamentous green algal species that occurs naturally in small amounts in Northern Michigan lakes. Its occurrence is governed by specific environmental requirements for temperature, substrate, nutrients, and other factors. It is found most commonly in the wave splash zone and shallow shoreline areas of lakes, as well as streams. It grows best on stable substrates such as rocks and logs, though artificial substrates such as

concrete or wood seawalls are also suitable. *Cladophora* prefers water temperatures in a range of 50 to 70 degrees Fahrenheit, which means that the optimal time for its growth and detection in Northern Michigan lakes is from mid-May to early July, and September to October.

The nutrients required for *Cladophora* to achieve large, dense growths are typically greater than the nutrient availability in the lakes of Northern Michigan. Therefore, shoreline locations where relatively high concentrations of nutrients, particularly phosphorus, are entering a lake can be identified by noting the presence of *Cladophora*. Although the growth features of *Cladophora* can be influenced by factors such as current patterns, shoreline topography, substrate composition, and wave action, the presence or absence of any significant growth is a powerful lake-wide screening tool. It can reveal the existence of chronic nutrient loading problems and assess the effectiveness of any remedial actions. Comparisons of the total number of algal growths can reveal trends in nutrient inputs due to changing land use.

Erosion along the shoreline has the potential to degrade the lake's water quality. Stormwater runoff through eroded areas carries sediments into the lake and impacts the lake ecosystem in a variety of ways. Sediments clog the gills of fish, aquatic insects, and other aquatic organisms. Excessive sediments smother fish spawning beds and fill interstitial spaces that provide habitat for a variety of aquatic organisms. Suspended sediments absorb sunlight energy and increase water temperatures. In addition, nutrients adhere to sediments that wash in from eroded areas, which can lead to nuisance aquatic plant growth and algal blooms.

Shoreline greenbelts are essential for maintaining a healthy aquatic ecosystem. A greenbelt consisting of a variety of native woody and herbaceous plant species provides habitat for near-shore aquatic organisms as well as other shoreline-dependent wildlife. They also help to stabilize shorelines against wave and ice action with their extensive network of deep, fibrous roots. Greenbelts also provide shade to nearshore areas, which is particularly important for lakes with cold water fisheries. In addition, greenbelts provide a mechanism to filter pollutants carried by stormwater from rain events and snowmelt.

Tributaries have a significant potential for influencing a lake's water quality as they are one of the primary conduits through which water is delivered to a lake from its watershed. Inlet streams may provide exceptionally high quality waters that benefit the lake ecosystem; conversely, they have the potential to deliver polluted waters that degrade the lake's water quality. Outlet streams flush water out of the lake, providing the means to remove contaminants that have accumulated in the lake ecosystem. With regard to shore surveys, noting the location of inlet tributaries is very helpful when evaluating shoreline algae conditions because nutrient concentrations are generally higher in streams than in lakes. The relatively higher nutrient levels delivered from streams often lead to naturally heavier *Cladophora* and other algal growth in nearby shoreline areas.

Lake-friendly shoreline property management is paramount for protecting water quality and sustaining a healthy, thriving lake ecosystem. Septic system maintenance, stormwater management, erosion control, and the elimination of fertilizers, herbicides, and pesticides are among the many low-cost best management practices that minimize the impact of shoreline properties on water quality.

### **Study Area:**

Intermediate Lake is located in the northwest Lower Peninsula of Michigan in Central Lake Township of Antrim County. Based on digitization of 2012 aerial orthophotography from the State of Michigan, the shoreline of Intermediate Lake measures 14.4 miles and lake surface area totals 1,570 acres. Intermediate Lake extends approximately six miles in a bent north to south direction and is between a tenth and a mile wide throughout its length. There are four distinguishable basins in Intermediate Lake; the north basin, mid-north basin, middle basin, and south basin. Both the north basin and mid-north basin are three quarter miles long and a third of a mile wide, the middle basin is three miles long and one mile wide, and the south basin is 1¼ miles long and half a mile wide (Figure 1). The basins are separated by three bottlenecks between a tenth and a quarter mile wide. Maps acquired from the Michigan Department of

Natural Resources (DNR) Institute for Fisheries Research indicate that the deepest point is located in the southeast side of the mid basin and measures approximately 70 feet.

The largest inlet is the Intermediate River, which flows into Intermediate Lake at the tip of the north basin, draining Hanley Lake and the remaining upper Elk River Chain of Lakes. The next largest tributaries are Openo and Fisk Creeks, which empty into the east side of the lake near the south end of the middle basin. The Intermediate River is the only outlet, which exits from the southwestern tip of the south basin.

According to GIS files developed by the Watershed Council using elevation data acquired from the State of Michigan, the Intermediate Lake Watershed encompasses approximately 72,258 acres, which includes the lake area (Figure 1). The immediate watershed accounts for 15,869 of these acres, while the remaining 56,389 acres are drainage through lakes upstream in the Chain. Land cover statistics were generated for the watershed using remote sensing data from the Coastal Great Lakes Land Cover project (NOAA 2010). Based on 2010 data, the majority of the watershed’s land-cover is natural; consisting primarily of forest, wetlands, and grassland (Table 1). There is a fair amount of agricultural land-cover in the watershed (26%) and some urban (3%). Both agricultural and urban land-cover changed by less than one percent between 1985 and 2010.

**Table 1.** Intermediate Lake Watershed land-cover statistics.

<b>Land Cover Type</b>	<b>1985 acres</b>	<b>1985 percent</b>	<b>2010 acres</b>	<b>2010 percent</b>	<b>Change (acres)</b>	<b>Change (percent)</b>
Agriculture	17916	24.8	18434	25.5	518	0.7
Barren	88	0.1	125	0.2	38	0.1
Forest	31019	42.9	30873	42.7	-146	-0.2
Grassland	4908	6.8	4193	5.8	-715	-1.0
Scrub/Shrub	1655	2.3	1847	2.6	192	0.3
Urban	2090	2.9	2163	3.0	73	0.1
Water	3113	4.3	3126	4.3	14	0.0
Wetland	11469	15.9	11495	15.9	26	0.0
<b>TOTAL</b>	<b>72258</b>	<b>100.0</b>	<b>72258</b>	<b>100.0</b>	<b>NA</b>	<b>NA</b>

The water quality of Intermediate Lake has been monitored consistently for two decades through the TOMWC's Volunteer Lake Monitoring program (VLM) and Comprehensive Water Quality Monitoring program (CWQM). TOMWC databases contain VLM and CWQM data that both date back to 1995. Data collected through these programs indicate good water quality.




Figure 1. Intermediate Lake: Features and Watershed.

Volunteer Lake Monitoring program data show that averaged Secchi disc depths on Intermediate Lake have ranged from 12 to 19 feet (Figure 2). Though there's not enough data to examine trends over time, water clarity is usually determined by two key factors: sediments and phytoplanktonic algae. Little sediment in the water is desirable, but too little algae can negatively impact the natural food web and lake ecosystem. A decrease in phytoplanktonic algae equates to a loss in primary productivity, which has cascading effects throughout the food chain. Invasive zebra mussels (*Dreissena polymorpha*), observed in Intermediate Lake during the survey, filter-feed on phytoplanktonic algae, which typically results in greater water clarity. Average chlorophyll-a measurements, which provide an indication of algal biomass in the lakes, typically decrease when zebra mussels become established in a lake; however, there isn't a large enough dataset to determine this trend (Figure 3).





Figure 2. Secchi disc depth data from Intermediate Lake (TOMWC 2016).




**Figure 3.** Chlorophyll-a data from Intermediate Lake (TOMWC 2016).

*\*Chlorophyll-a is measured in ug/L, which is micrograms per liter or parts per billion.*

Based on trophic status index data from the VLM program, Intermediate Lake borders between mesotrophy and oligotrophy. Oligotrophic lakes are typically large, deep, clear, and nutrient poor. In general, oligotrophic lakes contain high quality waters, but paradoxically have a lackluster fishery due to low biological productivity. Mesotrophic lakes are moderately productive. Total phosphorus data collected in the CWQM program show that levels have dropped throughout the last 20 years and are generally found below 10 parts per billion (PPB), which is typical for high quality lakes of Northern Michigan (Figure 4). This decrease is probably

due, at least in part, to the introduction of zebra mussels, which have filtered much of the algae out of the water column and disrupted the natural nutrient cycle in the lake.



**Figure 4.** Total phosphorus trends in Intermediate Lake 1995 to 2016 (TOMWC 2016).

*\*Total phosphorus is measured in ug/L, which is micrograms per liter or parts per billion.*

## METHODS

The Intermediate Lake shoreline was comprehensively surveyed in June of 2016 to document shoreline conditions that can potentially impact water quality. Shoreline conditions were surveyed by traveling in kayak as close to the shoreline as possible (usually within 20 feet) and noting *Cladophora* growth, substrate type, erosion, greenbelt health, shoreline alterations, and tributaries. A GPS camera was used to photograph all shoreline properties. Information for each property was recorded on field data sheets, subsequently inputted into a database, and used in conjunction with GPS data to link field data and photographs with property owner data from county equalization records.

### Field Survey Parameters

Shoreline property features were documented by photographing and noting physical features on a data sheet, such as building descriptions, public access sites, and county road endings. Due to data sheet space limits, building descriptions were recorded in an abbreviated style. For example, *Red 2 sty, brn rf, wht trm, fldstn chim, lg pine* means that the property has a red two-story house with a brown roof, white trim, fieldstone chimney, and a large pine tree in the yard. Whenever possible, names of property owners and addresses were included.

Developed parcels were noted on field data sheets and included as a separate column in the database. Properties described as developed indicate the presence of buildings or other significant permanent structures, including roadways, boat launching sites, and recreational properties (such as parks with pavilions and parking lots). Properties with only mowed or cleared areas, seasonal structures (such as docks or travel trailers), or unpaved pathways were not considered developed. Additionally, large parcels that had structures in an area far from the water's edge were not considered developed. The length and area of developed versus undeveloped shoreline was not calculated.

Many species of filamentous green algae are commonly found growing in the nearshore regions of lakes. Positive identification of these species usually requires the aid of a microscope.

However, *Cladophora* usually has an appearance and texture that is quite distinct to a trained surveyor, and these were the sole criteria upon which identification was based. Other species of filamentous green algae can respond to an external nutrient source in much the same way as *Cladophora*, though their value as an indicator species is not thought to be as reliable. When other species occurred in especially noticeable, large, dense growths, they were recorded on the data sheets and described the same as those of *Cladophora*.

When *Cladophora* was observed, it was described in terms of the length of shoreline with growth, the relative growth density, and any observed shoreline features potentially contributing to the growth. For example, “MHx30 – seeps” denotes a moderate to heavy growth that covered 30’ of the shoreline and with groundwater seeps in the area that may have been contributing to the growth. Both shoreline length and growth density are subjective estimates. Growth density is determined by estimating the percentage of substrate covered with *Cladophora* using the following categorization system:

**Table 2.** Categorization system for *Cladophora* density.

Density Category	Field Notation	Substrate Coverage
Very Light	(VL)	0% *
Light	(L)	1- 20%
Light to Moderate	(LM)	21-40%
Moderate	(M)	41-60%
Moderate to Heavy	(MH)	61-80%
Heavy	(H)	81-99%
Very Heavy	(VH)	90-100% *

*\*Very Light is noted when a green shimmer is noticed on hard substrate, but no filamentous growth present. Very Heavy overlaps with heavy and is distinguished by both high percentage of substrate coverage and long filamentous growth.*

Among other things, the distribution and size of each *Cladophora* growth is dependent on the amount of suitable substrate present. The extent of suitable substrate should therefore be

taken into account when interpreting the occurrence of individual growths, and assessing the overall distribution of *Cladophora* along a particular stretch of shoreline. Substrate types were noted during the survey, using the following abbreviations: m = soft muck or marl, s = sand, g = gravel (0.1" to 2.5" diameter), r = rock (2.5" to 10" diameter), b = boulder (>10" diameter), and w = woody debris. Substrate suitable for *Cladophora* growth include the g, r, b, and w types. The extent of suitable substrate along a shoreline parcel in terms of distance was not documented.

Erosion was noted based on shoreline areas that exhibited areas of bare soil, leaning or downed trees, exposed tree roots, undercut banks, slumping hunks of sod, or excessive deposits of sediments. Similar to *Cladophora*, shoreline erosion was recorded on field data sheets with estimates of its extent and relative severity (minor, moderate, or severe). For example "Mx20" indicated 20 feet of shoreline with moderate erosion. Additional information about the nature of the erosion, such as potential causes, was also noted.

Greenbelts (i.e., shoreline vegetation) were rated based on the length of shoreline with a greenbelt and the average depth of the greenbelt from the water's edge landward. Ratings for length ranged from 0 to 4, while ratings for depth ranged from 0 to 3. Ratings were based on the following:

**Length** ..... 0: None, 1: 1-10%, 2: 10-25%, 3: 25-75%, 4: >75%

**Depth** ..... 0: None, 1: <10 ft, 2: 10-40 ft, 3: >40 ft

Greenbelt ratings for length and depth were summed to produce an overall greenbelt score. Greenbelt scores ranged from 0 to 7, representing the greenbelt status or health. Scores of 0 were considered very poor, 1-2=poor, 3-4=moderate, 5-6=good, and 7=excellent.

Shoreline alterations were surveyed and noted with the following abbreviated descriptions: SB = steel bulkhead (i.e., seawall), BB = boulder bulkhead, CB = concrete bulkhead, RR = rock rip-rap, WB = wood bulkhead, BS = beach sand, BH = permanent boathouse, DP = discharge pipe.

Abbreviations were sometimes mixed or vary from what is listed above.

Tributaries (i.e., rivers and streams) were noted on the field data sheets and included in a separate column in the database. Additional information regarding shoreline property features or shoreline conditions recorded on field data sheets was included in the database in a “comments” column.

### **Data Processing**

Upon completing fieldwork, all field data were transferred to computer. Information from field data sheets was inputted into a Microsoft Excel® workbook. Digital photographs and GPS data were uploaded to a computer at the Watershed Council office and processed for use. Linking field and equalization data allows shoreline conditions documented during the survey to be referenced by parcel identification number or parcel owner name. Field data were linked to Antrim County parcel data in a GIS with the aid of GPS and photographs. Occasionally, errors occur wherein field data are not linked to the appropriate parcel.

In order to display survey results without pinpointing specific parcels, a new map layer was developed using the parcel map data layer acquired from the county equalization department and an Intermediate Lake shoreline layer. The new map layer consists of a narrow band following the shoreline, split into polygons that contain field and equalization data. This data layer was overlaid with other GIS data from the State of Michigan to produce a poster-size map to display survey results.

Final products include a comprehensive database, a complete set of GPS digital photographs, GIS data layers of shoreline parcels that include both county equalization and shore survey data, and a map displaying results. The database contains all data collected in the field and identification numbers in the database correspond to those in the GIS data layer and on hard-copy maps. GPS photographs were renamed using the same identification numbers and are linked to a GIS data layer.

## RESULTS

This survey documented shoreline conditions at 506 parcels on Intermediate Lake.

Approximately 86% (437) of shoreline properties on Intermediate Lake were considered to be developed.

Habitat generally considered suitable for *Cladophora* growth was present along at least part of the shoreline of 343 properties (68%). Noticeable growths of *Cladophora* or other filamentous green algae were found along the shoreline at 94 parcels (19% of the total or 27% of properties with suitable habitat). At properties where *Cladophora* growth was observed, nearly 80% consisted of light or very light growth, whereas only 8 parcels had growth in the moderate-heavy to heavy categories (Table 3).

**Table 3.** *Cladophora* density results.

<b><i>Cladophora</i> Density</b>	<b>Parcels</b>	<b>Percent</b>
Very light	41	8%
Light	34	7%
Light to Moderate	6	1%
Moderate	5	1%
Moderate to Heavy	0	0%
Heavy	6	1%
Very Heavy	2	0%
TOTAL	94	19%

*Cladophora* occurrence was more common in the northern portion of the Lake, compared to little growth within the southern portion (Figure 5). This shares a similar spatial distribution to patterns of development.




Figure 5. *Cladophora* algae density results for Intermediate Lake.

Greenbelt scores ranged from 0 (little to no greenbelt) to 7 (exemplary greenbelt). Less than one fourth of greenbelts (22%) along the Intermediate Lake shoreline were found to be in good or excellent condition (Table 4). The majority of parcels (63%) received a greenbelt rating in the poor or very poor categories.

**Table 4.** Greenbelt rating results.

<b>Greenbelt Rating</b>		<b>Number of Parcels</b>	<b>Percent of Parcels</b>
0	Very Poor (absent)	175	35%
1-2	Poor	144	28%
3-4	Moderate	75	15%
5-6	Good	52	10%
7	Excellent	59	12%

Poor greenbelts were clustered in the northern portion of Intermediate Lake, near the community of Central Lake and associated heavier development (Figure 6). Greenbelts improved to the south, but some clusters of development stood out as having little to no vegetative buffer at water's edge.




Figure 6. Greenbelt score totals results for Intermediate Lake.

Some form of shoreline alteration was noted at 77% of shoreline properties (Table 5). The majority of alterations consisted of boulder riprap (31%), while seawalls (wooden, concrete, or metal), were found at 13% of shoreline properties.

**Table 5.** Shoreline alteration results.

<b>Alteration Type</b>	<b>Number of Parcels*</b>	<b>Percent of Parcels With Alteration*</b>
Riprap (small)	125	25%
Riprap (boulder)	157	31%
Seawalls	66	13%
Beach Sand	123	24%
Unaltered	114	23%

*\*Numbers and percentages quantify alteration type, many parcels had multiple alterations*

Erosion was noted at 267 parcels (53%) on the Intermediate Lake shoreline (Table 6). 27% of shoreline properties with erosion were classified as minor in terms of severity, while roughly 3% of properties exhibited severe erosion.

**Table 6.** Shoreline erosion results.

<b>Erosion Category</b>	<b>Number of Properties</b>	<b>Percent of Properties*</b>
Minor	136	27%
Moderate	116	23%
Severe	15	3%
TOTAL	267	53%

Erosion was less common in the northern portions of Intermediate Lake. Mid-lake, especially along exposed shorelines at the Lake’s widest point, had erosion issues occurring more commonly (Figure 7).




Figure 7. Shoreline erosion severity results for Intermediate Lake.

## DISCUSSION

In general, development of shoreline parcels can negatively impact a lake's water quality due to a multitude of factors. Among the most serious impacts are: 1) loss of vegetation that would otherwise absorb and filter pollutants in stormwater runoff as well as stabilize shoreline areas and prevent erosion, 2) increased impervious surface area such as roofs, driveways and roads, which leads to greater inputs of stormwater runoff and associated pollutants, and 3) waste and byproducts of human activity such as septic leachate, fertilizers and decomposing yard waste that potentially reach and contaminate the lake water. Clearly, there are many problems associated with development, but there are also many solutions for reducing or even entirely eliminating impacts.

Numerous best management practices have been developed that help minimize negative impacts to water quality and which can be utilized during, or retroactively after, the development of shoreline parcels. A buffer of diverse, native plants can be maintained along the shoreline to filter pollutants and reduce erosion. Impacts from stormwater generated from roofs, roads, and driveways can be reduced using rain barrels, rain gardens, grassy swales, and many other techniques. Leachate reaching the lake from septic systems can be minimized by pumping the septic tank regularly, having all components of the septic system inspected regularly and replacing the septic system when necessary. Mulch can be composted far from the shoreline and fertilizers applied sparingly, if at all.

Results from the 2016 shoreline survey indicate that some of the aforementioned issues may pose a threat to the water quality and overall health of Intermediate Lake. Widespread removal of shoreline vegetation is the paramount concern. Over half (63%) of all shorelines exhibited greenbelts that were in poor condition. Erosion is also a concern, with light to moderate erosion commonly occurring throughout areas most exposed to wind and waves. Algal indicators of nutrient pollution are far less extensive than the above issues.

Comparisons with prior shoreline surveys show changes in these measurements over time. The

total number of properties with documented *Cladophora* growth increased by 13% since 2000 (Table 7). This indicates a slight trend of increasing near-shore nutrient pollution. Outreach regarding septic system maintenance, phasing out of old systems, and properly siting new systems may be playing a role in reduction in nutrient pollution related to septic systems. A few probable problem areas exist (as evidenced by heavy *Cladophora* growth occurring along the shoreline of eight parcels), which warrant further investigation. Where human-caused nutrient pollution is occurring, the source has to be identified in order to address the problem. Although impeded by factors such as wind, wave action, currents, and groundwater paths, efforts by trained personnel to identify specific nutrient input sources on individual properties are often successful.

**Table 7.** Critical shoreline survey parameter comparisons: 2000 to 2016.

<b>Survey Parameter</b>	<b>2000 Survey Results</b>		<b>2016 Survey Results</b>	
	Properties	%	Properties	%
<i>Cladophora</i> Algae Presence	31	6%	94	19%
Poor Greenbelts (score 0-2)	ND	ND	319	63%
Erosion	4	1%	267	53%
Shoreline Alterations	ND	ND	114	23%

The lack of vegetation on the Lake’s shoreline, which provides habitat and acts as a food source, can impact the abundance and diversity of aquatic organisms, ranging from minute crustaceans to top tier predator fish. Furthermore, the absence of vegetation leads to greater amounts of shoreline erosion and less filtration of pollutants. Shoreline survey data supports this relationship, with increases in erosion documented in tandem with declining greenbelt quality over the past years. Although a substantial number of greenbelts are in poor condition, 12% of properties received a perfect score, indicating exemplary greenbelt health. Properties with healthy, intact greenbelts provide a model for improvement for other shoreline properties.

**Table 8.** Shore survey statistics from Northern Michigan lakes.

Lake Name	Survey Date	<i>Cladophora</i> *	Heavy Algae*	Erosion*	Poor Greenbelts*	Alterations*
Beals Lake	2016	0%	0%	0%	17%	0%
Ben-Way Lake	2016	3%	0%	84%	47%	40%
Burt Lake	2009	47%	29%	4%	36%	46%
Bellaire Lake	2017	35%	1%	27%	30%	55%
Charlevoix, Lake	2012	22%	19%	14%	34%	79%
Clam Lake	2017	48%	5%	30%	51%	55%
Crooked Lake	2012	29%	26%	14%	51%	65%
Douglas Lake	2015	27%	6%	17%	53%	60%
Elk Lake	2017	84%	2%	52%	30%	87%
Ellsworth Lake	2016	40%	14%	38%	24%	23%
Hanley Lake	2016	11%	0%	33%	19%	23%
Huffman Lake	2015	14%	0%	7%	57%	70%
Huron, Duncan Bay	2013	41%	2%	19%	45%	63%
Huron, Grass Bay	2013	0%	0%	4%	0%	8%
Intermediate Lake	2016	19%	9%	53%	63%	77%
Lance Lake	2014	19%	0%	12%	35%	31%
Larks Lake	2006	4%	0%	ND	12%	29%
Mullett Lake	2016	44%	6%	36%	59%	76%
Pickerel Lake	2012	27%	33%	15%	52%	64%
Round Lake	2014	21%	0%	27%	44%	44%
Scotts Lake	2016	0%	0%	2%	18%	7%
Silver Lake	2014	3%	0%	70%	53%	65%
Skegemog Lake	2017	52%	5%	40%	46%	76%
St. Clair Lake	2016	4%	0%	13%	34%	21%
Six Mile Lake	2016	10%	24%	13%	41%	37%
Thayer Lake	2017	11%	1%	32%	16%	22%
Thumb Lake	2007	4%	0%	ND	ND	39%
Torch Lake	2017	39%	5%	26%	20%	ND
Walloon Lake	2016	62%	2%	17%	39%	80%
Wildwood Lake	2014	5%	0%	22%	45%	50%
Wilson	2016	27%	5%	29%	11%	14%
AVERAGE	NA	24%	6%	26%	36%	47%

*\*Percentages are in relation to number of parcels on the lake shore, except for “heavy algae”, which is the percent of only parcels that had Cladophora growth. Erosion is the percentage of parcels with moderate to severe erosion and poor greenbelts include those in the poor or very poor categories. ND=no data.*

Although many properties on Intermediate Lake are experiencing some form of erosion, the majority (51% of all erosion sites) are considered minor and only 6% of all erosion is considered to be severe. Many properties with patches of lawn at water's edge experience a minor undercutting caused by waves and ice shove. Properties with artificial beach sand usually experience some loss of sand into the lake, evidenced by small erosional rills leading into the lake. Although not catastrophic, these types of minor erosion do have the ability to degrade the water and habitat quality of Intermediate Lake. In addition, the wake from large boats may be exacerbating erosion as unnaturally large waves break along the shoreline.

The narrow northerly portion of the Lake limits water surface width for wind to generate waves (fetch), resulting in smaller naturally occurring waves and less regular impact to the shoreline from wave energy. Likewise, winter ice-shove is also less impactful in areas of lower fetch. Without exposure to erosional forces commonly found in areas of greater fetch, shorelines generally retain more vegetation and require less armoring. Furthermore, repair of the current erosion sites should be relatively easy to accomplish with plantings, as the low-energy shoreline allows for establishment of new plants. More exposed portions in the mid and southern portions of the Lake may pose a greater challenge for erosion site repair.

To prevent changes to the lake ecosystem, changes need to be made in shoreline property management. Mismanagement of shoreline properties can degrade the lake's water quality, diminish fisheries, and even create an environment that poses threats to human health.

## RECOMMENDATIONS

The full value of a shoreline survey is only achieved when the information is used to educate riparian property owners about preserving water quality, and to help them rectify any problem situations. The following are recommended follow-up actions:

1. Keep the specific results of the survey confidential (e.g., do not publish a list of sites where *Cladophora* algae were found) as some property owners may be sensitive to publicizing information regarding their property.
2. Send a general summary of the survey results to all shoreline residents, along with a packet of informational brochures produced by the Watershed Council and other organizations to provide information about threats to the Lake's ecosystem and public health as a result of poor shoreline property management practices as well as practical, feasible, and effective actions to protect water quality.
3. Organize and sponsor an informational session to present findings of the survey to shoreline residents and provide ideas and options for improving shoreline management practices that would help protect and improve the Lake's water quality.
4. Inform owners of properties with heavy *Cladophora* growths of specific results for their property, ask them to fill out a questionnaire in an attempt to interpret causes of the growth, and offer individualized recommendations for water quality protection. Following the questionnaire survey, property owners have the option to have the Watershed Council perform site visits and conduct groundwater testing in an effort to gain more insight into the nature of the findings. Again, it should be stressed that all information regarding names, specific locations, and findings be kept confidential to encourage property owner participation in this project.
5. Inform owners of properties with poor greenbelt scores and those with severely eroded shorelines of specific results for their property. Supply these property owners with information (e.g., brochures) regarding the benefits of greenbelts and/or the problems

associated with erosion. Encourage property owners to improve greenbelts using a mix of native plants and to correct erosion problems. Property owners can have the Watershed Council perform site assessments and carry out projects to improve greenbelts and/or correct erosion problems.

6. Utilize the Internet and other organizations' websites to share survey information. A general summary report and this detailed report can be posted websites because they do not contain any property-specific information. Property-specific information can be shared by randomizing and encrypting the shoreline survey database and providing property owners with a code number that refers specifically to survey results from their property. The Watershed Council is available to assist with this approach.
7. Continue to support the Tip of the Mitt Watershed Council Volunteer Lake and Stream Monitoring programs by providing volunteer support. The information collected by volunteers is extremely valuable for evaluating water quality and determining trends. Intermediate Lake Association is encouraged to continue supplying volunteer help and volunteers should attend training sessions held by the Watershed Council to ensure that a complete set of quality data is being collected each year.
8. Repeat some version of the survey periodically (ideally every 5 - 10 years), coupled with the follow-up activities described previously, in order to promote water quality awareness and good management practices on an ongoing basis. During each subsequent survey, more details about shoreline features are added to the database, which can be utilized for other water resource management applications.

## LITERATURE AND DATA REFERENCED

Antrim County Equalization/GIS. 2015. Antrim County GIS Parcel Data. Bellaire, MI. <http://www.antrimcounty.org/>

Michigan Department of Information Technology, Center for Geographic Information. 2015. Michigan Geographic Data Library. Lansing, MI. <http://www.mcgi.state.mi.us/mgdl/>

Michigan Department of Natural Resources. 2015. Inland Lake Maps. Lansing, MI. [http://www.michigan.gov/dnr/0,4570,7-153-10364\\_52261-67498--,00.html](http://www.michigan.gov/dnr/0,4570,7-153-10364_52261-67498--,00.html).

National Oceanic and Atmospheric Administration, Coastal Services Center. 2010. Coastal Great Lakes Land Cover Project. Charleston, SC. <http://coast.noaa.gov/dataregistry/search/collection/info/ccapregional>.

State of Michigan. 2012. Digital orthophotography from Michigan Imagery Solution Public Imagery. Lansing, MI. <http://mis.rsgis.msu.edu/mis/login.aspx>.

Tip of the Mitt Watershed Council. 2016. Comprehensive Water Quality Monitoring Program Data. Tip of the Mitt Watershed Council. Petoskey, MI. <http://www.watershedcouncil.org/Protect/>

Tip of the Mitt Watershed Council. 2016. Volunteer Lake Monitoring Program Data. Tip of the Mitt Watershed Council. Petoskey, MI. <http://www.watershedcouncil.org/Protect/>

Tip of the Mitt Watershed Council. 1998. A Shoreline Survey of Intermediate Lake. Tip of the Mitt Watershed Council. Petoskey, MI. <http://www.watershedcouncil.org/>